

STK24 Rabbit pAb

货号: **AYP15628**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:1000 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	47kDa/49kDa
实测分子量	49kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,A375,U-251MG,HT-29,NIH/3T3,Mouse liver,Mouse kidney,Mouse lung,Rat liver
细胞定位	Cytoplasm,Membrane,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 312-431 of human STK24 (NP_001027467.2).
序列	TDGQASGGSDSGDWIFTIREKDPKNLENGALQPSDLDRNKMKDIPKRPFSQLSTIISPLFAELKEKSQACGGNLSIEELRGAIYLAEAEACPGISDTMVAQLVQRLQRYSLSGGGTSSH

靶点信息

研究背景	This gene encodes a serine/threonine protein kinase that functions upstream of mitogen-activated protein kinase (MAPK) signaling. The encoded protein is cleaved into two chains by caspases; the N-terminal fragment (MST3/N) translocates to the nucleus and promotes programmed cells death. There is a pseudogene for this gene on chromosome X. Alternative splicing results in multiple transcript variants.
------	--

基因ID	8428
基因名	STK24
Swiss	Q9Y6E0
别名	STK24;HEL-S-95;MST3;MST3B;STE20;STK3

产品验证



Western blot analysis of STK24 expressed in Mouse liver,Rat liver,HeLa using STK24 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blotting buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn