

— ABLYBIO, Help Your Research



Phospho-MEF2C-S396 Rabbit pAb

货号: **AYP15652**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	44kDa/47kDa/50kDa/51kDa/52kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562,MCF7
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	A synthetic phosphorylated peptide around S396 of human MEF2C (NP_002388.2).
------	--

靶点信息

研究背景	This locus encodes a member of the MADS box transcription enhancer factor 2 (MEF2) family of proteins, which play a role in myogenesis. The encoded protein, MEF2 polypeptide C, has both trans-activating and DNA binding activities. This protein may play a role in maintaining the differentiated state of muscle cells. Mutations and deletions at this locus have been associated with severe mental retardation, stereotypic movements, epilepsy, and cerebral malformation. Alternatively spliced transcript variants have been described.
基因ID	4208
基因名	MEF2C
Swiss	Q06413 (https://www.uniprot.org/uniprotkb/Q06413/entry)
别名	MEF2C,C5DELq14.3,DEL5q14.3,Phospho-MEF2C-S396 Rabbit pAb,Myocyte enhancer factor 2C

产品验证



Western blot analysis of Phospho-MEF2C-S396 expressed in K-562, MCF7 using Phospho-MEF2C-S396 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)