

REPIN1 Rabbit pAb

货号: **AYP15654**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	
实测分子量	64kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562,HeLa,Mouse testis,Mouse liver
细胞定位	nucleoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 376-567 of human REPI N1 (NP_037532.2).
序列	SCDDCGRSFRLERFLRAHQHQHTGERPFTCAECGKNFGKKTHLVAHSRVHSGERPFACEECGRRFSQGSHLAAHRRDH APDRPFVCPDCGKAFRHKPYLAAHRRHTGEKPYVCPDCGKAFSQKSNLVSHRRIHTGERPYACPDCDRSFSQKSNLITHR KSHIRDGAFCCAICGQTFDDEERLLAHQKKHDV

靶点信息

研究背景	Sequence-specific double-stranded DNA-binding protein required for initiation of chromosomal DNA replication. Binds on 5'-ATT-3' reiterated sequences downstream of the origin of bidirectional replication (OBR and a second, homologous ATT sequence of opposite orientation situated within the OBR zone. Facilitates DNA bending.
基因ID	29803

基因名	REPIN1
Swiss	Q9BWE0
别名	AP4;RIP60;ZNF464;Zfp464;REPIN1

产品验证



Western blot analysis of REPIN1 expressed in Mouse testis, K-562, HeLa using REPIN1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn