

UQCC2 Rabbit pAb

货号: **AYP15677**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	14kDa
实测分子量	15kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,THP-1,HepG2,U-251MG,Mouse heart,Mouse brain,Mouse testis,Rat brain
细胞定位	Mitochondrion,Mitochondrion inner membrane,Mitochondrion intermembrane space,Mitochondrion matrix ,mitochondrion nucleoid
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 14-126 of human UQCC 2 (NP_115716.1).
序列	EEWPVDETKRGRDLGAYLRQRVAQAFREGENTQVAEPEACDQMYESLARLHSNYYKHKYPRPRDTSFSGLSLEEYKLILSTDTLEELKEIDKGMWKKLQEKFAPKGPEEDHKA

靶点信息

研究背景	This gene encodes a nucleoid protein localized to the mitochondria inner membrane. The encoded protein affects regulation of insulin secretion, mitochondrial ATP production, and myogenesis through modulation of mitochondrial respiratory chain activity.
------	--

基因ID	84300
基因名	UQCC2
Swiss	Q9BRT2
别名	UQCC2;C6orf125;C6orf126;Cbp6;M19;MNF1;bA6B20.2

产品验证



Western blot analysis of UQCC2 expressed in Mouse heart,Rat brain,MCF7 using UQCC2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane . Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn