

— ABLYBIO, Help Your Research



# EIF3L Rabbit pAb

货号: **AYP15682**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:1000 - 1:4000 <b>IHC:</b> 1:50 - 1:200
理论分子量	61kDa/66kDa
实测分子量	67kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,HT-1080,NIH/3T3,Mouse heart,Mouse pancreas,Mouse liver
细胞定位	Cytoplasm
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-230 of human EIF3L (NP_057175.1).
------	---

## 靶点信息

研究背景	Component of the eukaryotic translation initiation factor 3 (eIF-3 complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA <sup>i</sup> and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression.
基因ID	51386
基因名	EIF3L
Swiss	Q9Y262 ( <a href="https://www.uniprot.org/uniprotkb/Q9Y262/entry">https://www.uniprot.org/uniprotkb/Q9Y262/entry</a> )
别名	EIF3L,EIF3EIP,EIF3S11,EIF3S6IP,HSPC021,HSPC025,MSTP005,EIF3L Rabbit pAb,Eukaryotic translation initiation factor 3 subunit 6-interacting protein,Eukaryotic translation initiation factor 3 subunit E-interacting protein

## 产品验证



Western blot analysis of EIF3L expressed in Mouse heart, HepG2, HT-1080 using EIF3L Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))