

TRIP12 Rabbit pAb

货号: **AYP15692**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:100 IF/ICC: 1:50 - 1:100
理论分子量	192kDa/220kDa/223kDa/225kDa
实测分子量	260kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat
细胞定位	Nucleus,nucleoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-240 of human TRIP12 (NP_001271143.1).
序列	MSNRPNNNPGGSLRRSQRNTAGAQPDQDDSIGGRSHLGQAKHKGYSPPESRKSNSKAPKVQSNTTSELGRHLSKRSCS SSSAVIVQPDPDRANTSERQKTGQVPPKDNSRGVKSASPDYRNRTNSPSSAKKPKALQHTESPSETNKPHSKSKRHL DQEQLKSAQSPSTSKAHRKSGATGGRSRQKRKRTESSCVKS GSGSESTGAEERSAKPTKLASKSATSAGCSTITDS SS

靶点信息

研究背景	The protein encoded by this gene is an E3 ubiquitin-protein ligase involved in the degradation of the p19A RF/ARF isoform of CDKN2A, a tumor suppressor. The encoded protein also plays a role in the DNA damage response by regulating the stability of USP7, which regulates tumor suppressor p53.
------	--

基因ID	9320
基因名	TRIP12
Swiss	Q14669
别名	TRIP12;TRIP-12;ULF

产品验证



Western blot analysis of TRIP12 expressed in HeLa, Jurkat using TRIP12 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5 % non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn