

CD72 Rabbit pAb

货号: **AYP15708**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	40kDa
实测分子量	40KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Daudi
细胞定位	Membrane,Single-pass type II membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 120-359 of human CD72 (NP_001773.1).
序列	QVSQQLQQTNRVLEVTNSSLRQQLRLKITQLGQSAEDLQGSRRRELQSQEALQVEQRAHQAAEGQLQACQADRQKTKE TLQSEEQRRALEQKLSNMENRLKPFFTCGSADTCCPSGWIMHQKSCFYISLTSKNWQESQKQCETLSSKLATFSEIYPQ SHSYFLNSLLPNGGSGNSYWTGLSSNKDWKLTDDTQRTRTYAQSSKCNKVHKTWSWWTLESESCRSSLPYICEMTAF RFPD

靶点信息

研究背景	B-Cell Differentiation Antigen CD72 (CD72) is a single-pass type II membrane protein. CD72 exists as a disulfide-linked homodimer and contains one C-type lectin domain. CD72 is expressed on B lineage cells, N K cells, monocytes, dendritic cells, and mast cells. CD72 is a ligand for CD5. CD72 associates with CD5, in teracts with PTPN6/SHP-1 and plays a role in B-cell proliferation and differentiation. CD72 associates with CD79A in the B cell antigen receptor (BCR) complex following antigen stimulation and dampens BCR signaling through interactions with the phosphatase SHP-1.
------	--

基因ID	971
基因名	CD72
Swiss	P21854
别名	CD72;CD72b;LYB2

产品验证



Western blot analysis of CD72 expressed in Daudi using CD72 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn