

SLC39A2 Rabbit pAb

货号: **AYP15712**

产品信息

反应	Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	
实测分子量	33KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat brain
细胞定位	cytoplasmic ribonucleoprotein granule,cytoplasmic vesicle,plasma membrane
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human SLC39A2 (NP_055394.2).
序列	MEQLLGIKLGCLFALLALTLGCGLTPICFKWFQIDAARGHRLVLRLLGCISAGVFLGAGFMHMTAEALEEIESQIQKFMVQ NRSASERNSSGDADSAHM

靶点信息

研究背景	This gene encodes a member of the ZIP family of metal ion transporters. The encoded protein functions as a zinc transporter. Mutations in this gene may be associated with susceptibility to carotid artery disease. Multiple transcript variants have been described. [provided by RefSeq, Mar 2010]
基因ID	29986

基因名	SLC39A2
Swiss	Q9NP94
别名	6A1;ETI-1;ZIP-2;ZIP2;SLC39A2

产品验证



Western blot analysis of SLC39A2 expressed in Rat brain using SLC39A2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn