

— ABLYBIO, Help Your Research



# TXNDC9 Rabbit pAb

货号: **AYP15772**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:100
理论分子量	22kDa/26kDa
实测分子量	26kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,293T,HeLa,Jurkat,Mouse testis,Mouse heart
细胞定位	Cytoplasm,Midbody,Nucleus,centrosome,cytoskeleton,microtubule organizing center
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-110 of human TXNDC 9 (NP_005774.2).
------	---

## 靶点信息

研究背景	The protein encoded by this gene is a member of the thioredoxin family. The exact function of this protein is not known but it is associated with cell differentiation.
基因ID	10190
基因名	TXNDC9
Swiss	O14530 ( <a href="https://www.uniprot.org/uniprotkb/O14530/entry">https://www.uniprot.org/uniprotkb/O14530/entry</a> )
别名	TXNDC9,APACD,PHLP3,TXNDC9 Rabbit pAb,ATP-binding protein associated with cell differentiation,Protein 1-4

## 产品验证



Western blot analysis of TXNDC9 expressed in Mouse testis,U-87MG,293T using TXNDC9 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))