

TXNDC9 Rabbit pAb

货号: **AYP15772**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:100
理论分子量	22kDa/26kDa
实测分子量	26kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,293T,HeLa,Jurkat,Mouse testis,Mouse heart
细胞定位	Cytoplasm,Midbody,Nucleus,centrosome,cytoskeleton,microtubule organizing center
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-110 of human TXNDC 9 (NP_005774.2).
序列	MEADASVDMFSKVLEHQLLQTTKLVEEHL DSEIQKLDQMD EDELERLKEKRLQALRKAQQKQEWLSKGHGEYREIPSE R DFFQEVKESENVVCHF YRDSTFRCKILDRH

靶点信息

研究背景	The protein encoded by this gene is a member of the thioredoxin family. The exact function of this protein is not known but it is associated with cell differentiation.
基因ID	10190

基因名	TXNDC9
Swiss	O14530
别名	TXNDC9;APACD;PHLP3

产品验证



Western blot analysis of TXNDC9 expressed in Mouse testis,U-87MG,293T using TXNDC9 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn