

— ABLYBIO, Help Your Research



# ZIC3 Rabbit pAb

货号: **AYP15798**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200
理论分子量	50kDa
实测分子量	51kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,Mouse brain,Rat brain
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-240 of human ZIC3 (N P_003404.1).
------	---

## 靶点信息

研究背景	This gene encodes a member of the ZIC family of C2H2-type zinc finger proteins. This nuclear protein probably functions as a transcription factor in early stages of left-right body axis formation. Mutations in this gene cause X-linked visceral heterotaxy, which includes congenital heart disease and left-right axis defects in organs.
基因ID	7547
基因名	ZIC3
Swiss	O60481 ( <a href="https://www.uniprot.org/uniprotkb/O60481/entry">https://www.uniprot.org/uniprotkb/O60481/entry</a> )
别名	ZIC3,HTX,HTX1,VACTERLX,ZNF203,ZIC3 Rabbit pAb,Zinc finger protein 203,Zinc finger protein of the cerebellum 3

## 产品验证



Western blot analysis of ZIC3 expressed in MCF7, Mouse brain, Rat brain using ZIC3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)