

# ITIH3 Rabbit pAb

货号: **AYP15800**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	99kDa
实测分子量	105kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,293T,NCI-H460,Mouse liver,Mouse lung,Rat liver
细胞定位	Secreted
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 35-180 of human ITIH3 (NP_002208.3).
序列	LPEGVANGIEVYSTKINSKVTSRFAHNVVMTMRVNRADTAKEVSFDVELPKTAFITNFTLTIDGVTPGNVKEKEVAKKQYEKAVSQGKTAGLVKASGRKLEKFTVSVNVAAGSKVTFELTYEELLKRHKGYEMYLKVQPKQLVK

## 靶点信息

研究背景	This gene encodes the heavy chain subunit of the pre-alpha-trypsin inhibitor complex. This complex may stabilize the extracellular matrix through its ability to bind hyaluronic acid. Polymorphisms of this gene may be associated with increased risk for schizophrenia and major depressive disorder. This gene is present in an inter-alpha-trypsin inhibitor family gene cluster on chromosome 3.
基因ID	3699

基因名	ITIH3
Swiss	Q06033
别名	ITIH3;H3P;ITI-HC3;SHAP

## 产品验证



Western blot analysis of ITIH3 expressed in Mouse liver,Rat liver,HepG2 using ITIH3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)