

CDCA2 Rabbit pAb

货号: **AYP15817**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	7kDa/112kDa
实测分子量	130kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	HeLa, Jurkat
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 724-1023 of human CD CA2 (NP_689775.2).
序列	SDSPKALTLQQGQEFSSAGGQNAENLCQFFKISPDNLNICKERKDDFLGAAEGKLQCNRLMPNSQKDCHCLGDVLIENTKE SKSQSEDLGRKPMESSVSVSCRDRKDRRRSMCYSDGRSLHLEKNGNHTPSSVSGSSVEISLENSELFKDLSDAIEQTFQR RNSETKVRRSTRLQKDLENEGLVWISLPLPSTSQKAKRRITICTFDSSGFESMSPIKETVSSRQKPMAPPVSDPENSQGP AGSSDEPGKRRKSFICISTLANTKATSQFKGYRRRSSLNKGESSLTALERIEHNGERKQ

靶点信息

研究背景	This gene encodes a targeting subunit of the cell-cycle associated protein, protein phosphatase 1, with a role in targeting this protein to chromatin during anaphase. These two proteins comprise a phosphatase c complex that is involved in nuclear envelope reformation and regulation of the DNA damage response. The encoded protein may also play a role in cancer progression. Alternative splicing results in multiple transcript variants.
------	--

基因ID	157313
基因名	CDCA2
Swiss	Q69YH5
别名	CDCA2;PPP1R81;Repo-Man

产品验证



Western blot analysis of CDCA2 expressed in HeLa, Jurkat using CDCA2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5 % non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn