

# TRMT6 Rabbit pAb

货号: **AYP15827**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	36kDa/40kDa/44kDa/55kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG, HepG2
细胞定位	Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human TRMT6 (NP_057023.2).
序列	MEGSGEQPGPQPQHPGDHRIRDGDFVVLKREDVFKAVQVQRRKKVTFEKQWFYLDNVIGHSYGTAFEVTSGGSLQPKK KREPTAETKEAGTDNRNIVDDGKSQKLTQDDIKALKDKGKIGEEIVQQLIENSTTFRDKTEFAQDKYIKKKKKYEAITVV KPSTRILSIMYYAREPGKINHMRDYDTLAQMLTLGNIRAG

## 靶点信息

研究背景	This gene encodes a member of the tRNA methyltransferase 6 protein family. A similar protein in yeast is part of a two component methyltransferase, which is involved in the posttranslational modification that produces the modified nucleoside 1-methyladenosine in tRNAs. Modified 1-methyladenosine influences initiator methionine stability and may be involved in the replication of human immunodeficiency virus type 1. Alternative splicing results in multiple transcript variants.
------	---

基因ID	51605
基因名	TRMT6
Swiss	Q9UJA5
别名	TRMT6;CGI-09;GCD10;Gcd10p

## 产品验证



Western blot analysis of TRMT6 expressed in U-87MG,HepG2 using TRMT6 Rabbit pAb at 1:1000. Second ary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer : 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)