

NUP160 Rabbit pAb

货号: **AYP15837**

产品信息

| | |
|-------|--|
| 反应 | Human |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB IHC |
| 推荐浓度 | WB: 1:500 - 1:1000 IHC: 1:20 - 1:50 |
| 理论分子量 | 24kDa/55kDa/162kDa |
| 实测分子量 | 160kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HeLa,A-431,Jurkat,HepG2 |
| 细胞定位 | Nucleus,nuclear pore complex |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 40-150 of human NUP160 (NP_056046.1). |
| 序列 | ALERSFVELSGAERERPRHFREFTVCSIGTANAVAGAVKYESAGGFYYVESGKLFVTRNRFIHWKTSGDTLELMEESLDI NLLNNAIRLKFQNCVLPGGVYVSETQNR |

靶点信息

| | |
|------|--|
| 研究背景 | NUP160 is 1 of up to 60 proteins that make up the 120-MD nuclear pore complex, which mediates nucleoplasmic transport.[supplied by OMIM, Apr 2004] |
| 基因ID | 23279 |

| | |
|-------|--------|
| 基因名 | NUP160 |
| Swiss | Q12769 |
| 别名 | NUP160 |

产品验证



Western blot analysis of NUP160 expressed in HeLa,A-431,Jurkat using NUP160 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn