

— ABLYBIO, Help Your Research



NADK Rabbit pAb

货号: **AYP15841**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:1000 - 1:2000
理论分子量	46kDa/49kDa/63kDa
实测分子量	49kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Mouse liver,Mouse brain,Rat liver,Rat brain
细胞定位	cytosol
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-180 of human NADK (NP_075394.3).
------	--

靶点信息

研究背景	NADK catalyzes the transfer of a phosphate group from ATP to NAD to generate NADP, which in its reduced form acts as an electron donor for biosynthetic reactions (Lerner et al., 2001 [PubMed 11594753]).[supplied by OMIM, Mar 2008]
基因ID	65220
基因名	NADK
Swiss	O95544 (https://www.uniprot.org/uniprotkb/O95544/entry)
别名	NADK,dj283E3.1,NADK Rabbit pAb,Poly(P)/ATP NAD kinase

产品验证



Western blot analysis of NADK expressed in Mouse liver,Rat liver,293T using NADK Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)