

EPB41L2 Rabbit pAb

货号: **AYP15845**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:200 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	76kDa/83kDa/95kDa/112kDa
实测分子量	180kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,HeLa,Jurkat,SKOV3
细胞定位	Cell membrane,Cytoplasm,cytoskeleton
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-210 of human EPB41L2 (NP_001186317.1).
序列	MTTEVGSVSEVKKDSSQLGTDATKEKPKEVAENQQNQSSDPPEEEKGSQPPPAAESQSSLRQKREKETSESRGISRFIPPWLKKQKSYTLVVAKDGGDKKEPTQAVVEEQVLDKEEPLPEEQRQAKGDAEEMAQKKQEIKVEVKEEKPSVSKEEKPSVSKVEMQPTELVSKEREKVKETQEDKLEGGAAKRETKEVQTNELKAEKASQK

靶点信息

研究背景	Required for dynein-dynactin complex and NUMA1 recruitment at the mitotic cell cortex during anaphase .
基因ID	2037

基因名	EPB41L2
Swiss	O43491
别名	EPB41L2;4.1-G;4.1G

产品验证



Western blot analysis of EPB41L2 expressed in U-87MG, HeLa, Jurkat using EPB41L2 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn