

DDT Rabbit pAb

货号: **AYP15881**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:200 - 1:2000
理论分子量	11kDa/12kDa
实测分子量	13kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,A-549,Mouse liver,Rat liver
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-118 of human DDT (NP_001346.1).
序列	MPFLELDTNLPANRVPAGLEKRLCAAASILGKPADRVNVTVRPGLAMALSGSTEPCAQLSISISIGVVGTAEDNRSHSAHF FEFLTKELALGQDRILIRFFPLESWQIGKIGTVMFTL

靶点信息

研究背景	D-dopachrome tautomerase converts D-dopachrome into 5,6-dihydroxyindole. The DDT gene is related to the migration inhibitory factor (MIF) in terms of sequence, enzyme activity, and gene structure. DDT and MIF are closely linked on chromosome 22.
基因ID	1652

基因名	DDT
Swiss	P30046
别名	DDT;DDCT

产品验证



Western blot analysis of DDT expressed in Mouse liver, Rat liver, 293T using DDT Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn