

PRAP1 Rabbit pAb

货号: **AYP15889**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	10kDa/16kDa/17kDa
实测分子量	17kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	293T, HeLa, LO2
细胞定位	Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-151 of human PRAP1 (NP_660203.3).
序列	MRRLLLVTSLVVLLWEAGAVPAPKVPKMQVKHWPSEQDPEKAWGARVVEPPEKDDQLVVLFPVQKPKLLTTEEKPRGQGRGPILPGTKAWMETEDTLGHVLSPEPDHDSLHPPPEEDQGEERPRLWVMPNHQVLLGPEEDQDHIYHPQ

靶点信息

研究背景	Lipid-binding protein which promotes lipid absorption by facilitating MTTP-mediated lipid transfer (mainly triglycerides and phospholipids) and MTTP-mediated apoB lipoprotein assembly and secretion (By similarity). Protects the gastrointestinal epithelium from irradiation-induced apoptosis (By similarity). May play an important role in maintaining normal growth homeostasis in epithelial cells. Involved in p53/TP53-dependent cell survival after DNA damage. May downregulate the expression of MAD1L1 and exert a suppressive role in mitotic spindle assembly checkpoint in hepatocellular carcinomas.
------	---

基因ID	118471
基因名	PRAP1
Swiss	Q96NZ9
别名	PRAP1;PRO1195;UPA

产品验证



Western blot analysis of PRAP1 expressed in 293T, HeLa, LO2 using PRAP1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn