

— ABLYBIO, Help Your Research



GTF2E2 Rabbit pAb

货号: **AYP15897**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:100
理论分子量	33kDa
实测分子量	33KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,Mouse testis
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-291 of human GTF2E2 (NP_002086.1).
------	--------------------------------------------------------------------------------------------------------------------

靶点信息

研究背景	The general transcription factor IIE (TFIIE) is part of the RNA polymerase II transcription initiation complex, recruiting TFIIF and being essential for promoter clearance by RNA polymerase II. TFIIE is a heterodimer (and sometimes heterotetramer) of alpha and beta subunits. The protein encoded by this gene represents the beta subunit of TFIIE. [provided by RefSeq, Jan 2017]
基因ID	2961
基因名	GTF2E2
Swiss	P29084 (https://www.uniprot.org/uniprotkb/P29084/entry)
别名	GTF2E2,FE,TF2E2,TFIIE-B,TTD6,GTF2E2 Rabbit pAb,General transcription factor IIE subunit 2

产品验证



Western blot analysis of GTF2E2 expressed in Raji, Mouse testis using GTF2E2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)