

# HFM1 Rabbit pAb

货号: **AYP15917**

## 产品信息

|       |  |
|-------|--|
| 反应    | Human,Mouse,Rat  |
| 宿主    | Rabbit   |
| 克隆性   | Polyclonal   |
| 预测反应  |  |
| 应用    | WB IHC   |
| 推荐浓度  | <b>WB:</b> 1:500 - 1:2000<br><b>IHC:</b> 1:50 - 1:100  |
| 理论分子量 | 62kDa/162kDa   |
| 实测分子量 | 130kDa   |
| 形式    | Liquid   |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物   | Unconjugated   |
| 阳性对照  | K-562  |
| 细胞定位  | nucleus  |
| 纯化    | Affinity purification  |

## 抗原信息


|      |   |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 1-240 of human HFM1 ( NP_001017975.4).  |
| 序列   | MLKSNDCLFSLLENLFFPEKDEVENHPDNEKSLDWFLPPAPLISEIPDTQELEEELESHKLLGQEKRPKMLTSNLKITNEDTNY<br>ISLTQKFQFAFPSDKYEQDDLNLEGVGNNDLSHVAGKLTYSQKYKNHIGTEIAPEKSVPDDTKLVNFAEDKGESTSVFRK<br>RLFKISDNIHGSAYSNDNELDSHIGSVKIVQTEMNKGKSRNYSNSKQKFQYSANVFTANNAFSAEIGEGMFKAPS |

## 靶点信息

|      |   |
|------|---|
| 研究背景 | The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). |
| 基因ID | 164045  |

|       |   |
|-------|---|
| 基因名   | HFM1  |
| Swiss | A2PYH4  |
| 别名    | HFM1;MER3;POF9;SEC63D1;Si-11;Si-11-6;helicase |

## 产品验证



Western blot analysis of HFM1 expressed in K-562 using HFM1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)