

C17orf70 Rabbit pAb

货号: **AYP15924**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	27kDa/77kDa/93kDa
实测分子量	110kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,A-431,HeLa,Mouse brain,Mouse liver,Rat brain
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 652-881 of human C17orf70 (NP_079437.5).
序列	CLRFPGGLAPPHTRAPSPGLGPTDRDPVATFLETCPREPGSQPAGPASLRAEYLPSPVASIKVSAELLRAALKDGHSGVPLCCATLQWLLAENAAVDVVRARALSSIQGVAPDGANVHLIVREVAMTDLCPAGPIQAVEIQVESSSLADICRAHHAVVGRMQTMVTEQATQGSSAPDLRVQYLRQIHANHETLLREVQTLRDRLCTEDEASSCATAQRLLQVYRQLRHPSLILL

靶点信息

研究背景	FAAP100 is a component of the Fanconi anemia (FA; MIM 277650) core complex and is required for core complex stability and FANCD2 (see MIM 227646) monoubiquitination (Ling et al., 2007 [PubMed 17396147]).[supplied by OMIM, Mar 2008]
基因ID	80233

基因名	FAAP100
Swiss	Q0VG06
别名	FAAP100;C17orf70

产品验证



Western blot analysis of C17orf70 expressed in Mouse brain,Rat brain,SH-SY5Y using C17orf70 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 12 0s.

实验步骤

访问官网浏览详情: www.ablybio.cn