

# FCGRT Rabbit pAb

货号: **AYP15942**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:1000 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	39kDa
实测分子量	40kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HL-60
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 27-290 of human FcRn ( NP_004098.1).
序列	HLSLLYHLTAVSSPAPGTPAFWVSWGWLGPQQYLSYNLSRGEAEPCGAWVWENQVSWYWEKETTDLRIKEKLFLEAFKALGGKGPYTLQGLLGCELGPDNTSVPTAKFALNGEEFMNFDLKQGTWGGDWPEALAIQRWQQQDKAANKELTFLLFSCP HRLREHLERGRGNLEWKEPPSMRLKARSSPGFSVLTCSAFSFYPPQLRFLRNLGLAAGTGQGDGFPNSDGSFHASSS LTVKSGDEHHYCCIVQHAGLAQPLRVEL

## 靶点信息

研究背景	This gene encodes a receptor that binds the Fc region of monomeric immunoglobulin G. The encoded protein transfers immunoglobulin G antibodies from mother to fetus across the placenta. This protein also binds immunoglobulin G to protect the antibody from degradation. Alternative splicing results in multiple transcript variants.
------	---

基因ID	2217
基因名	FCGRT
Swiss	P55899
别名	FCGRT;FCRN;alpha-chain

## 产品验证



Western blot analysis of FCGRT expressed in HL-60 using FCGRT Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)