

— ABLYBIO, Help Your Research



NAA25 Rabbit pAb

货号: **AYP15956**

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	99kDa/112kDa
实测分子量	120kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse uterus,Mouse eye,Mouse ovary,Rat uterus,Rat eye,Rat ovary
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 703-972 of human NAA 25 (NP_079229.2).
------	--

靶点信息

研究背景	This gene encodes the auxiliary subunit of the heteromeric N-terminal acetyltransferase B complex. This complex acetylates methionine residues that are followed by acidic or asparagine residues.
------	--

基因ID	80018
基因名	NAA25
Swiss	Q14CX7 (https://www.uniprot.org/uniprotkb/Q14CX7/entry)
别名	NAA25,C12orf30,MDM20,NAP1,NAA25 Rabbit pAb,Mitochondrial distribution and morphology protein 20,N-terminal acetyltransferase B complex subunit MDM20,N-terminal acetyltransferase B complex subunit NAA25,p120

产品验证



Western blot analysis of NAA25 expressed in Mouse uterus,Rat uterus,Mouse eye using NAA25 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)