

RPP30 Rabbit pAb

货号: **AYP15975**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:100 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	29kDa/35kDa
实测分子量	29kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,HT-1080,HepG2
细胞定位	Nucleus,nucleolus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-268 of human RPP30 (NP_006404.1).
序列	MAVFADLDLRAGSDLKALRGLVETA AHLGYSVVAINHIVDFKEKKQEIEKPVAVSELTTLPIVQGKSRPIKILTRLTIIVSDPS HCNVL RATSSRARLYDVVAVFPKTEKLFHIACTHLDVLDVCITVTEKLPFYFKRPPINVAIDRGLAFELVYSPA IKDSTMRRYTI SSALNLMQICKGKNVISSAAERPLEIRGPYDVANLGLLFGLS ESDAKAAVSTNCRAALLHGETRKTAFGIISTVKKPRPSEG DEDCLPASKKAKCEG

靶点信息

研究背景	Component of ribonuclease P, a ribonucleoprotein complex that generates mature tRNA molecules by cleaving their 5'-ends. Also a component of the MRP ribonuclease complex, which cleaves pre-rRNA sequences.
------	--

基因ID	10556
基因名	RPP30
Swiss	P78346
别名	RPP30;TSG15

产品验证



Western blot analysis of RPP30 expressed in HeLa, Jurkat, HT-1080 using RPP30 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn