

# ZBTB17 Rabbit pAb

货号: **AYP15980**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:100 - 1:200
理论分子量	79kDa/87kDa/88kDa
实测分子量	103kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	B cells,U-87MG,Hela
细胞定位	Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 121-360 of human ZBTB 17 (NP_003434.2).
序列	PGGNAEALATEGGDKRAKEEKVATSTLSRLEQAGRSTPIGPSRDLKEERGGQAQSAASGAEQTEKADAPREPPPVELKPD PTSGMAAAEAEAALSESSEQEMEVEPARKGEEEQKEEQEEEGAGPAEVKEEGSQLENGEAPEENENEESAGTDSGQE LGSEARGLRSGTYGDRTESKAYGSVIHKCEDCGKEFTHTGNFKRHIRIHTGEKPFSCRECSKAFSDPAACKAHEKTHSPLK

## 靶点信息

研究背景	This gene encodes a zinc finger protein involved in the regulation of c-myc. The symbol MIZ1 has also been associated with PIAS2 which is a different gene located on chromosome 18.
基因ID	7709

基因名	ZBTB17
Swiss	Q13105
别名	ZBTB17;MIZ-1;ZNF151;ZNF60;pHZ-67

## 产品验证



Western blot analysis of ZBTB17 expressed in B cells,U-87MG,HeLa using ZBTB17 Rabbit pAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blo cking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)