

NCOR2 Rabbit pAb

货号: **AYP16009**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:100
理论分子量	81kDa/268kDa/273kDa/274kDa
实测分子量	275kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,LO2
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 670-900 of human NCO R2 (NP_006303.4).
序列	KLKMEKERNARRKKKKAPAAASEEAAFPVVEDEEMEASGVSGNEEEMVEEAEALHASGNEVPRGECSPATVNNSSDT ESIPSPHTEAAKDTGQNGPKPPATLGADGPPPGPTPPPEDIPAPTEPTPASEATGAPTPPPAPPSPSAPPVVPKKEEKEET AAAPPVEEGEEQPPAAEELAVDTGKAEPPVKSECTEEAEEGPAKKGKDAEAAEATAEGALKAEKKEGGS

靶点信息

研究背景	This gene encodes a nuclear receptor co-repressor that mediates transcriptional silencing of certain target genes. The encoded protein is a member of a family of thyroid hormone- and retinoic acid receptor-associated co-repressors. This protein acts as part of a multisubunit complex which includes histone deacetylases to modify chromatin structure that prevents basal transcriptional activity of target genes. Aberrant expression of this gene is associated with certain cancers. Alternate splicing results in multiple transcript variants encoding different isoforms.
基因ID	9612
基因名	NCOR2
Swiss	Q9Y618
别名	NCOR2;CTG26;N-CoR2;SMAP270;SMRT;SMRTE;SMRTE-tau;TNRC14;TRAC;TRAC-1;TRAC1

产品验证



Western blot analysis of NCOR2 expressed in HeLa, LO2 using NCOR2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn