

TAMM41 Rabbit pAb

货号: **AYP16013**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	35kDa/38kDa/51kDa
实测分子量	38kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,K-562,Jurkat
细胞定位	Matrix side,Mitochondrion inner membrane,Peripheral membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 25-316 of human TAMM 41 (NP_620162.1).
序列	SLAFVYGSGVYRQAGPSSDQKNAMLDVFVTDVDDPVAWHSKLNKKNWSHYSFLKVLGPKIITSIQNNYAGVYYNSLIMC NGRLIKYGVISTNVLIEDLLNWNLYIAGRLQKPKIISVNEVDVTLRSALDRNLKSAVTA AFLMLPESFSEEDLFIEIAGLSYSG DFRMVVGEDKTKVLNIVKPNIAHFRELYGSILQENPQVVYKSQQGWLEIDKSPEGQFTQLMTLPKTLQQQINHIMDPPGKN RDVEETLFQVAHDPDCGDVVRGLKKSVIYSSKLKHKMWKGWLRKTS

靶点信息

研究背景	Catalyzes the conversion of phosphatidic acid (PA to CDP-diacylglycerol (CDP-DAG, an essential intermediate in the synthesis of phosphatidylglycerol, cardiolipin and phosphatidylinositol.
基因ID	132001

基因名	TAMM41
Swiss	Q96BW9
别名	TAMM41;C3orf31;RAM41;TAM41

产品验证



Western blot analysis of TAMM41 expressed in MCF7, K-562, Jurkat using TAMM41 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn