

# PGLS Rabbit pAb

货号: **AYP16017**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200
理论分子量	27kDa
实测分子量	28kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW620,THP-1,HepG2,HeLa,U-251MG,Mouse heart,Mouse spleen,Mouse liver,Rat brain
细胞定位	Cytoplasm
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-258 of human PGLS ( NP_036220.1).
序列	MAAPAPGLISVVFSSSQELGAALQLVAQRAACCLAGARARFALGLSGGSLVSMMLARELPAAVAPAGPASLARWTLGFCDE RLVPFDHAESTYGLYRTHLLSRLPIPESQVITINPELPVEEAAEDYAKKLRQAFQGDSIPVFDLLILGVGPDGHTCSLFPDHPL LQEREKIVAPISDSPKPPQVRVTLTPVLNAARTVIFVATGEGKAAVLKRILEDQEENPLPAALVQPHTGKLCWFLDEAAARLL TVPFKHOSTL

## 靶点信息

研究背景	Hydrolysis of 6-phosphogluconolactone to 6-phosphogluconate.
基因ID	25796

基因名	PGLS
Swiss	O95336
别名	PGLS;6PGL;HEL-S-304

## 产品验证



Western blot analysis of PGLS expressed in Mouse heart,Rat brain,SW620 using PGLS Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)