

# RNF13 Rabbit pAb

货号: **AYP16020**

## 产品信息

反应	Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	29kDa/42kDa
实测分子量	47kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver,Mouse kidney
细胞定位	Endoplasmic reticulum membrane,Golgi apparatus membrane,Late endosome membrane,Lysosome membrane,Nucleus inner membrane,Single-pass membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 204-381 of human RNF13 (NP_009213.1).
序列	TKFVQDRHRARRNRLRKDQLKLPVHKFKKGDEYDVCAICLDEYEDGDKLRILPCSHAYHCKCVDWPWLTKTKTCPVCKQKVVPSQGDSDSDTDSSQEENEVTEHTPLLRPLASVSAQSFGALSESRSHQNMTESSDYEDDNETDSSDAENEINEHDVVVQLQPNGERDYNIANTV

## 靶点信息

研究背景	The protein encoded by this gene contains a RING zinc finger, a motif known to be involved in protein-protein interactions. The specific function of this gene has not yet been determined. Alternatively spliced transcript variants that encode the same protein have been reported. A pseudogene, which is also located on chromosome 3, has been defined for this gene.
------	---

基因ID	11342
基因名	RNF13
Swiss	O43567
别名	RNF13;RZF

## 产品验证



Western blot analysis of RNF13 expressed in Mouse liver, Mouse kidney using RNF13 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)