

— ABLYBIO, Help Your Research



MID2 Rabbit pAb

货号: **AYP16035**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	79kDa/83kDa
实测分子量	83kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BT-474,293T,U-251MG,Mouse lung
细胞定位	Cytoplasm,cytoskeleton
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 476-705 of human MID2 (NP_438112.2).
------	--

靶点信息

研究背景	The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. The protein localizes to microtubular structures in the cytoplasm. Alternate splicing of this gene results in two transcript variants encoding different isoforms.
基因ID	11043
基因名	MID2
Swiss	Q9UJV3 (https://www.uniprot.org/uniprotkb/Q9UJV3/entry)
别名	MID2,FXY2,MRX101,RNF60,TRIM1,midline 2,MID2 Rabbit pAb,Midin-2,Midline defect 2,Midline-2,RING finger protein 60,RING-type E3 ubiquitin transferase MID2,Tripartite motif-containing protein 1

产品验证



Western blot analysis of MID2 expressed in Mouse lung, BT-474, 293T using MID2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)