

— ABLYBIO, Help Your Research



# ULBP2 Rabbit pAb

货号: **AYP16051**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:100 - 1:500
理论分子量	27kDa
实测分子量	35KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	THP-1
细胞定位	Cell membrane,GPI-anchor,Lipid-anchor,Secreted
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 26-216 of human ULBP2 (NP_079493.1).
------	--

## 靶点信息

研究背景	This gene encodes a major histocompatibility complex (MHC) class I-related molecule that binds to the NKG2D receptor on natural killer (NK) cells to trigger release of multiple cytokines and chemokines that in turn contribute to the recruitment and activation of NK cells. The encoded protein undergoes further processing to generate the mature protein that is either anchored to membrane via a glycosylphosphatidylinositol moiety, or secreted. Many malignant cells secrete the encoded protein to evade immunosurveillance by NK cells. This gene is located in a cluster of multiple MHC class I-related genes on chromosome 6.
基因ID	80328
基因名	ULBP2
Swiss	Q9BZM5 ( <a href="https://www.uniprot.org/uniprotkb/Q9BZM5/entry">https://www.uniprot.org/uniprotkb/Q9BZM5/entry</a> )
别名	ULBP2,ALCAN-alpha,N2DL2,NKG2DL2,RAET1H,ULBP2 Rabbit pAb,NKG2D ligand 2,Retinoic acid early transcript 1H

## 产品验证



Western blot analysis of ULBP2 expressed in THP-1 using ULBP2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)