

BAIAP2L1 Rabbit pAb

货号: **AYP16058**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	56kDa
实测分子量	57kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,HeLa,293T,A-549,Mouse lung
细胞定位	Cytoplasm,cytoskeleton
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 198-388 of human BAIA P2L1 (NP_061330.2).
序列	DKHCGFANHIHYHLQSAELLNSKLPRWQETCVDIAIKVPEKIMNMIEEIKTPASTPVSQTPQASPMIERSNVVRKDYDTLSK CSPKMPPAPSGRAYTSPLIDMFNNPATAAPNSQQRVNNSTGTSQSLQSVSVATGLNMMKKQKVKTIFFHTAGSNKTLT SFAQGDVITLLIPEEKDQWLYGEHDVSKA

靶点信息

研究背景	This gene encodes a member of the IMD (IRSp53/MIM homology domain) family. Members of this family can be subdivided in two groups, the IRSp53-like and MIM-like, based on the presence or absence of the SH3 (Src homology 3) domain. The protein encoded by this gene contains a conserved IMD, also known as F-actin bundling domain, at the N-terminus, and a canonical SH3 domain near the C-terminus, so it belongs to the IRSp53-like group. This protein is the substrate for insulin receptor tyrosine kinase and binds to the small GTPase Rac. It is involved in signal transduction pathways that link deformation of the plasma membrane and remodeling of the actin cytoskeleton. It also promotes actin assembly and membrane protrusions when overexpressed in mammalian cells, and is essential to the formation of a potent actin assembly complex during EHEC (Enterohemorrhagic Escherichia coli) pedestal formation.
基因ID	55971
基因名	BAIAP2L1
Swiss	Q9UHR4
别名	BAIAP2L1;IRTKS

产品验证



Western blot analysis of BAIAP2L1 expressed in Mouse lung, HepG2, HeLa using BAIAP2L1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn