

— ABLYBIO, Help Your Research



STK38 Rabbit pAb

货号: **AYP16068**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	54kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,SW480,293T,MCF7,Mouse liver
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human STK38 (NP_009202.1).
------	-------------------------------------------------------------------------------------------------------------------

靶点信息

研究背景	This gene encodes a member of the AGC serine/threonine kinase family of proteins. The kinase activity of this protein is regulated by autophosphorylation and phosphorylation by other upstream kinases. This protein has been shown to function in the cell cycle and apoptosis. This protein has also been found to regulate the protein stability and transcriptional activity of the MYC oncogene. Alternative splicing results in multiple transcript variants.
基因ID	11329
基因名	STK38
Swiss	Q15208 (https://www.uniprot.org/uniprotkb/Q15208/entry)
别名	STK38,NDR,NDR1,STK38 Rabbit pAb,NDR1 protein kinase,Nuclear Dbf2-related kinase 1

产品验证



Western blot analysis of STK38 expressed in Mouse liver,A-549,SW480 using STK38 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)