

FBXW4 Rabbit pAb

货号: **AYP16074**

产品信息

反应	Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	46kDa
实测分子量	46kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain
细胞定位	cytosol
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 143-412 of human FBXW4 (NP_071322.1).
序列	QFRPDGASLNRRPLGVFAGHDEDVCHFVLANSRSHVSAAGDGGKIGIHKIHSTFTVKYSAHEQEVNVCVDCCKGGIIVSGSRDR TAKVWPLASGRLGQCLHTIQTEDRVWSIAISPLLSSFVTGTACCGHFSPLRIWDLNSGQLMTHLGSDFPPGAGVLDVMEY SPFTLLSCGYDITYVRYWDLRTSVRKCVMEWEEPHDSTLYCLQTDGNHLLATGSSYYGVVRLWDRRQRACLHAFPLTSTP LSSPVYCLRLTTKHLYAALSYNLHVLDLDFQNP

靶点信息

研究背景	This gene is a member of the F-box/WD-40 gene family, which recruit specific target proteins through their WD-40 protein-protein binding domains for ubiquitin mediated degradation. In mouse, a highly similar protein is thought to be responsible for maintaining the apical ectodermal ridge of developing limb buds; disruption of the mouse gene results in the absence of central digits, underdeveloped or absent metacarpal/metatarsal bones and syndactyly. This phenotype is remarkably similar to split hand-split foot malformation in humans, a clinically heterogeneous condition with a variety of modes of transmission. An autosomal recessive form has been mapped to the chromosomal region where this gene is located, and complex rearrangements involving duplications of this gene and others have been associated with the condition. A pseudogene of this locus has been mapped to one of the introns of the BCR gene on chromosome 22.
基因ID	6468
基因名	FBXW4
Swiss	P57775
别名	FBXW4;DAC;FBW4;FBWD4;SHFM3;SHSF3

产品验证



Western blot analysis of FBXW4 expressed in Mouse brain using FBXW4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn