

MonoMethyl-NONO/p54nrb-R357/R365 Rabbit pAb

货号: **AYP16091**

产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:100 IF/ICC: 1:50 - 1:100
理论分子量	43kDa/54kDa
实测分子量	43kDa/54kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Nucleus,Nucleus speckle,nucleolus
纯化	Affinity purification

抗原信息


抗原信息	A synthetic monomethylated peptide around R357 & R365 of human NONO/p54nrb (Q15233).
序列	RQQEEMRR

靶点信息

研究背景	This gene encodes an RNA-binding protein which plays various roles in the nucleus, including transcriptional regulation and RNA splicing. A rearrangement between this gene and the transcription factor E3 gene has been observed in papillary renal cell carcinoma. Alternatively spliced transcript variants have been described. Pseudogenes exist on Chromosomes 2 and 16.
------	---

基因ID	4841
基因名	NONO
Swiss	Q15233
别名	NONO;MRXS34;NMT55;NRB54;P54;P54NRB;PPP1R114

产品验证



Western blot analysis of MonoMethyl-NONO/p54nrb-R357/R365 expressed in human liver cancer, r at testis using MonoMethyl-NONO/p54nrb-R357/R365 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn