

SSR4 Rabbit pAb

货号: **AYP16113**

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:200 - 1:2000 |
| 理论分子量 | 18kDa |
| 实测分子量 | 19kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | LO2,U-87MG,MCF7,HL-60,Mouse liver,Mouse brain |
| 细胞定位 | Endoplasmic reticulum membrane,Single-pass type I membrane protein |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 24-144 of human SSR4 (NP_006271.1). |
| 序列 | EACLEPQITPSYYTTS DAVISTETVFIVEISLTCKNRVQNMALYADVGGKQFPVTRGQDVGRYQVSWSLDHKSAHAGTYEV RFFDEESYLLRKAQRNEDISIIPPLFTVSDHRGTWNG |

靶点信息

| | |
|------|--|
| 研究背景 | This gene encodes the delta subunit of the translocon-associated protein complex which is involved in translocating proteins across the endoplasmic reticulum membrane. The encoded protein is located in the Xq28 region and is arranged in a compact head-to-head manner with the isocitrate dehydrogenase 3 (NAD+) gamma gene and both genes are driven by a CpG-embedded bidirectional promoter. Alternate splicing results in multiple transcript variants. |
| 基因ID | 6748 |

| | |
|-------|------------------|
| 基因名 | SSR4 |
| Swiss | P51571 |
| 别名 | SSR4;CDG1Y;TRAPD |

产品验证



Western blot analysis of SSR4 expressed in Mouse liver, LO2, U-87MG using SSR4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn