

SIK3 Rabbit pAb

货号: **AYP16116**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	65kDa/125kDa/133kDa/139kDa
实测分子量	145KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293F
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 60-310 of human SIK3 (NP_079440.3).
序列	PARIGYYEIDRTIGKGNFAVVKRATHLVTKAKVAIKIIDKTQLDEENLKKIFREVQIMKMLCHPHIIRLYQVMETERMIYLVTEYA SGGEIFDHLVAHGRMAEKEARRKFKQIVTAVYFCHCRNIVHRDLKAENLLLDANLNIAKIDFGFSNLFTPGQLLKTWCGSPP YAAPELFEGKEYDGPVKVDIWSLGVVLYLVCGALPFDGSTLQNLRRARVLSGKFRIPFFMSTECEHLIRHMLVLDPNKRLSME Q

靶点信息

研究背景	Enables ATP binding activity; magnesium ion binding activity; and protein serine/threonine kinase activity . Involved in positive regulation of TORC1 signaling; positive regulation of TORC2 signaling; and protein p hosphorylation. Located in nucleoplasm.
基因ID	23387

基因名	SIK3
Swiss	Q9Y2K2
别名	SIK3;L19;QSK;SIK-3

产品验证



Western blot analysis of SIK3 expressed in 293F using SIK3 Rabbit pAb at 1:1000. Secondary antibody: HR P Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn