

# B9D1 Rabbit pAb

货号: **AYP16132**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	16kDa/22kDa
实测分子量	23kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Mouse testis,Mouse lung,Mouse ovary,Mouse kidney
细胞定位	Cytoplasm,cilium axoneme,cilium basal body,cytoskeleton
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-130 of human B9D1 ( NP_056496.1).
序列	MATASPSVFLMVGQVESAQFPEYDDLYCKYCFVYGQDWAPTAGLEEGISQITSKSDVRQALVWNFPIDVTFKSTNPY GWPQIVLSVYGPDPVFGNDVVRGYGAVHVPFSPGRHKRTIPMFVPESTSKL

## 靶点信息

研究背景	This gene encodes a B9 domain-containing protein, one of several that are involved in ciliogenesis. Alterations in expression of this gene have been found in a family with Meckel syndrome. Meckel syndrome has been associated with at least six different genes. This gene is located within the Smith-Magenis syndrome region on chromosome 17.
------	---

基因ID	27077
基因名	B9D1
Swiss	Q9UPM9
别名	B9D1;B9;EPPB9;JBTS27;MKS9;MKS1

## 产品验证



Western blot analysis of B9D1 expressed in Mouse testis, HeLa, Mouse lung using B9D1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)