

NPTN Rabbit pAb

货号: **AYP16134**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	30kDa/31kDa/37kDa/43kDa/44kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SKOV3,Jurkat
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 29-200 of human NPTN (NP_059429.1).
序列	QNEPRIVTSEEVIIRDSPVLPVTLQC�LTSSSHTLTYSYWTKNGVELSATRKNASNMEYRINKPRAEDSGEYHCVYHFVSAP KANATIEVKAAPDITGHKRSENKNEGQDATMYCKSVGYPHDPWIWRKKENGMPMDIVNTSGRFFIINKENYTELNIVNLQI TEDPGEYEC

靶点信息

研究背景	This gene encodes a type I transmembrane protein belonging to the Ig superfamily. The protein is believed to be involved in cell-cell interactions or cell-substrate interactions. Alternative splicing results in multiple transcript variants.
基因ID	27020

基因名	NPTN
Swiss	Q9Y639
别名	NPTN;GP55;GP65;SDFR1;SDR1;np55;np65

产品验证



Western blot analysis of NPTN expressed in SKOV3, Jurkat using NPTN Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn