

# CHIC2 Rabbit pAb

货号: **AYP16135**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:100
理论分子量	19kDa
实测分子量	19kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse testis
细胞定位	Cell membrane,Cytoplasmic vesicle
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-165 of human CHIC2 ( NP_036242.1).
序列	MADFDEIYEEEEDEERALEEQLLKYPDPVVVRGSGHVTVFGLSNKFESEFPSSLTGKVAPEEFKASINRVNSCLKKNLPVN VRWLLCGCLCCCCTLGCSMWPVICLSKRTRRSIEKLEWENNRLYHKLCLHWRLSKRKCETNNMMEYVILIEFLPKTPIFRPD

## 靶点信息

研究背景	This gene encodes a member of the CHIC family of proteins. The encoded protein contains a cysteine-rich hydrophobic (CHIC) motif, and is localized to vesicular structures and the plasma membrane. This gene is associated with some cases of acute myeloid leukemia.
基因ID	26511

基因名	CHIC2
Swiss	Q9UKJ5
别名	CHIC2;BTL

## 产品验证



Western blot analysis of CHIC2 expressed in Mouse testis using CHIC2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)