

ACAT2 Rabbit pAb

货号: **AYP16183**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	41kDa/44kDa
实测分子量	41kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BT-474,MCF7,HeLa,HepG2,PC-12,Mouse liver
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 150-250 of human ACAT2 (NP_005882.2).
序列	GLTDAFHNMHGITAENVAKKWQVSREDQDKVAVLSQNRTEAQAQKAGHFDKEIVPVLVSTRKGLIEVKTDEFPRHGSNIEAMSKLKPFLDTGTGTPAN

靶点信息

研究背景	The product of this gene is an enzyme involved in lipid metabolism, and it encodes cytosolic acetoacetyl-CoA thiolase. This gene shows complementary overlapping with the 3-prime region of the TCP1 gene in both mouse and human. These genes are encoded on opposite strands of DNA, as well as in opposite transcriptional orientation. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
------	---

基因ID	39
基因名	ACAT2
Swiss	Q9BWD1
别名	ACAT2

产品验证



Western blot analysis of ACAT2 expressed in Mouse liver, BT-474, MCF7 using ACAT2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn