

ADAR1 Rabbit pAb

货号: **AYP16184**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:100 - 1:500 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	103kDa/131kDa/133kDa/136kDa/140kDa
实测分子量	110KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293F
细胞定位	Cytoplasm,Nucleus,Nucleus,nucleolus
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 150-250 of human ADAR11 (NP_001102.2).
序列	LGEGKATTAHDLGKLGTPKKEINRVLYSLAKKGKLGKEAGTPPLWKIAVSTQAWNQHSGVVRPDGHSQGAPNSDPSLEPEDRNSTSVSEDLLEPFIAVSA

靶点信息

研究背景	This gene encodes the enzyme responsible for RNA editing by site-specific deamination of adenosines. This enzyme destabilizes double-stranded RNA through conversion of adenosine to inosine. Mutations in this gene have been associated with dyschromatosis symmetrica hereditaria. Alternative splicing results in multiple transcript variants.
------	---

基因ID	103
基因名	ADAR
Swiss	P55265
别名	ADAR;ADAR1;AGS6;DRADA;DSH;DSRAD;G1P1;IFI-4;IFI4;K88DSRBP;P136

产品验证



Western blot analysis of ADAR1 expressed in 293F using ADAR1 Rabbit pAb at 1:1000. Secondary antibody y: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn