

— ABLYBIO, Help Your Research



ADAR1 Rabbit pAb

货号: **AYP16184**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:100 - 1:500 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	103kDa/131kDa/133kDa/136kDa/140kDa
实测分子量	110KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293F
细胞定位	Cytoplasm,Nucleus,Nucleus,nucleolus
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 150-250 of human ADAR11 (NP_001102.2).
------	---

靶点信息

研究背景	This gene encodes the enzyme responsible for RNA editing by site-specific deamination of adenosines. This enzyme destabilizes double-stranded RNA through conversion of adenosine to inosine. Mutations in this gene have been associated with dyschromatosis symmetrica hereditaria. Alternative splicing results in multiple transcript variants.
基因ID	103
基因名	ADAR
Swiss	P55265 (https://www.uniprot.org/uniprotkb/P55265/entry)
别名	ADAR,ADAR1,AGS6,DRADA,DSH,DSRAD,G1P1,IFI-4,IFI4,K88DSRBP,P136,ADAR1 Rabbit pAb,136 kDa double-stranded RNA-binding protein,Interferon-inducible protein 4

产品验证



Western blot analysis of ADAR1 expressed in 293F using ADAR1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)