

# ALG2 Rabbit pAb

货号: **AYP16188**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	37kDa/47kDa
实测分子量	47kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,Mouse brain
细胞定位	Membrane,Single-pass membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 197-416 of human ALG2 (NP_149078.1).
序列	DVLYPSLNVTSFDSVPEKLDLVLVPGKGFLLLSINRYERKKNLTLALEALVQLRGRLLTSQDWERVHLIVAGGYDERVLENV EHYQELKKMVQSDLGQYVTFLLRSFSDKQKISLLHSCCTCVLYTPSNEHFGIVPLEAMYMCPVIAVNSGGPLESIDHSVTG FLCEPDPVHFSEAIEKFIREPSLKATMGLAGRARVKEKFSPEAFTEQLYRYVTKLLV

## 靶点信息

研究背景	This gene encodes a member of the glycosyltransferase 1 family. The encoded protein acts as an alpha 1, 3 mannosyltransferase, mannosylating Man(2)GlcNAc(2)-dolichol diphosphate and Man(1)GlcNAc(2)-dolichol diphosphate to form Man(3)GlcNAc(2)-dolichol diphosphate. Defects in this gene have been associated with congenital disorder of glycosylation type 1h (CDG-1h). Alternative splicing results in multiple transcript variants.
------	--

基因ID	85365
基因名	ALG2
Swiss	Q9H553
别名	ALG2;CDG1I;CDGII;CMS14;CMSTA3;NET38;hALPG2;alpha-1;3/1

## 产品验证



Western blot analysis of ALG2 expressed in MCF7, Mouse brain using ALG2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)