

PRDM6 Rabbit pAb

货号: **AYP16191**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	14kDa/46kDa/64kDa
实测分子量	80kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,MCF7,Mouse lung
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 376-595 of human PRDM6 (NP_001129711.1).
序列	LQCIAQDENLNPSTVMEAMCRQDALQPFNKSSKLAPTTQRSVVFQTPCSRNFSLLDKSGPIESGFNQINVKNQRVLA SPTSTSQLHSEFSDWHLWKCGQCFKFTTQRILLQMHVCTQNPDRPYQCGHCSQSFQSPSELRNHVVTHSSDRPFKCGY CGRAFAGATLNNHIRTHTGEKPFKCERCERSFTQATQLSRHQRMPECKPITESPESIEVD

靶点信息

研究背景	Putative histone methyltransferase that acts as a transcriptional repressor of smooth muscle gene expression. Promotes the transition from differentiated to proliferative smooth muscle by suppressing differentiation and maintaining the proliferative potential of vascular smooth muscle cells. Also plays a role in endothelial cells by inhibiting endothelial cell proliferation, survival and differentiation. It is unclear whether it has histone methyltransferase activity in vivo. According to some authors, it does not act as a histone methyltransferase by itself and represses transcription by recruiting EHMT2/G9a. According to others, it possesses histone methyltransferase activity when associated with other proteins and specifically methylates 'Lys-20' of histone H4 in vitro. 'Lys-20' methylation represents a specific tag for epigenetic transcriptional repression.
基因ID	93166
基因名	PRDM6
Swiss	Q9NQX0
别名	PRDM6;KMT8C;PDA3;PRISM

产品验证



Western blot analysis of PRDM6 expressed in Jurkat, MCF7, Mouse lung using PRDM6 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn