

RAB17 Rabbit pAb

货号: **AYP16195**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	9kDa/23kDa
实测分子量	23kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A375,HepG2,22Rv1,SW480
细胞定位	Cell projection,Cytoplasmic side,Lipid-anchor,Recycling endosome membrane,dendrite
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-212 of human RAB17 (NP_071894.1).
序列	MAQAHRTQPRAAPSQPRVFKLVLLGSGSVGKSSALALRYVKNDFKSILPTVGCAFFTKVVDVGATSLKLEIWDTAGQEKY HSVCHLYFRGANAALLVYDITRKDSFLKAQQWLKDLEELHPGEVLVMLVGNKTDLSQEREVTFQEGKEFADSQKLLFME TSAKLNHQVSEVFNTVAQELLQRSDEEGQALRGDAVAVALNKGPARQAKCCAH

靶点信息

研究背景	The Rab subfamily of small GTPases plays an important role in the regulation of membrane trafficking. RAB17 is an epithelial cell-specific GTPase (Lutcke et al., 1993 [PubMed 8486736]).[supplied by OMIM, Oct 2009]
基因ID	64284

基因名	RAB17
Swiss	Q9H0T7
别名	RAB17

产品验证



Western blot analysis of RAB17 expressed in A375, HepG2, 22Rv1 using RAB17 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn