

# BSND Rabbit pAb

货号: **AYP16216**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	35kDa
实测分子量	35kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Rat liver,Rat kidney,Rat intestine
细胞定位	Cell membrane,Cytoplasm,Multi-pass membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 54-320 of human BSND (NP_476517.1).
序列	CQCYPKITFVPADSDFQGILSPKAMGLENGLAEMKSPSPQPPYVRLWEEAAYDQSLPDFSHIQMKVMSYSEDHRSLLA PEMGQPKLGTSDGGEGGPGDVQAWMEAAVVIHKGSDSEGERRLTQSWPGPLACPQGPAPLASFQDDLDMDSSEGS SPNASPHDREEACSPQQEPQGCRCLDRFQDFALIDAPLTDLEPQEGQQWEIALPNNWQRYPRTKVEEKEASDTGGEEP EKEEEDLYGLPDGAGDLLPKELGFEPDTQG

## 靶点信息

研究背景	This gene encodes an essential beta subunit for CLC chloride channels. These heteromeric channels localize to basolateral membranes of renal tubules and of potassium-secreting epithelia of the inner ear. Mutations in this gene have been associated with Bartter syndrome with sensorineural deafness.
基因ID	7809

基因名	BSND
Swiss	Q8WZ55
别名	BSND;BART;DFNB73;barttin

## 产品验证



Western blot analysis of BSND expressed in Rat liver,293T,Rat kidney using BSND Rabbit pAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)