

TTF2 Rabbit pAb

货号: **AYP16219**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	75kDa/129kDa
实测分子量	130kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	MCF7, HL-60, SKOV3, Jurkat
细胞定位	Cytoplasm, Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-270 of human TTF2 (NP_003585.3).
序列	MEEVRCPEHGTFCFLKTGVRDGPNGKGSFYVCRADTCSFVRATDIPVSHCLLHEDFVVELQGLLLPQDKKEYRLFFRCIRS KAEGRWCGSIPWQDPDSKEHSVSNKSHASETFHHSSNWLRNPFKVLDPKQEPALWKQLIKGEGEEKKADKKQREK GDQLFDQKKEQKPEMMEKDLSSGLVPKPKKQSVVQEKQEEGAEIQCEAETGGTHKRDSEIKSQCQGNELTRPSASS QEKSSGKSQDVQRESEPLREKVTQLLPQNVHSHN

靶点信息

研究背景	This gene encodes a member of the SWI2/SNF2 family of proteins, which play a critical role in altering protein-DNA interactions. The encoded protein has been shown to have dsDNA-dependent ATPase activity and RNA polymerase II termination activity. This protein interacts with cell division cycle 5-like, associates with human splicing complexes, and plays a role in pre-mRNA splicing.
------	--

基因ID	8458
基因名	TTF2
Swiss	Q9UNY4
别名	TTF2;HuF2;ZGRF6

产品验证



Western blot analysis of TTF2 expressed in MCF7,HL-60,SKOV3 using TTF2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn