

AAR2 Rabbit pAb

货号: **AYP16263**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	
实测分子量	43kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,A-549
细胞定位	
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-250 of human AAR2 (NP_056326.2).
序列	MAAVQMDPELAKRLLFFEGATVVILNMPKGTTEFGIDYNSWEVGPKFRGVKMIPPGIHFHLYSSVDKANPKEVGPRMGFFLS LHQRLTVLRWSTLREEVDLSPAPESEVEAMRANLQELDQFLGPPYATLKKWISLTNFISEATVEKLQPENRQICAFSDVL PVL SMKHTKDRVGQNLPRCGIECKSYQEGLARLPEMKPRAGTEIRFSELPTQMFPEGATPAEITKHSMDLSYALETVLNKQF PSSPQD

靶点信息

研究背景	This gene encodes the homolog of the yeast A1-alpha2 repressin protein that is involved in mRNA splicing . Alternately spliced transcript variants have been found for this gene. [provided by RefSeq, Dec 2012]
基因ID	25980

基因名	AAR2
Swiss	Q9Y312
别名	C20orf4;CGI-23;AAR2

产品验证



Western blot analysis of AAR2 expressed in MCF7, A-549 using AAR2 Rabbit pAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn